

PRODUCT INSERT

Nutri-Fly™

German Food “Sick Fly” Formulation

Powdered Drosophila Media by Genesee Scientific

Description: Nutri-Fly German Food “Sick Fly” Formulation is made to accommodate the requirements of large, communal Drosophila media preparation facilities as well as small, individual laboratories. Recipe can be doubled, tripled, etc. in the same batches to meet any media quantity requirements. Sealed in single-use foil pouches, Nutri-Fly GF is not only convenient but also robust, and guaranteed to make fresh and substantive food throughout its shelf-life of one year from delivery. Strict lot traceability procedures are employed to each ingredient, in every batch.

Directions: Empty entire contents of packet into a cooking vessel. Add $\frac{1}{2}$ - $\frac{3}{4}$ specified RT H₂O and stir out any clumps. Add remaining RT H₂O and stir. Bring to a boil, then immediately turn off heat source. Allow food to cool to 60°C to 40°C. Add specified (Table 1.) propionic acid^{††} quantity (not supplied) and pour.

Ingredients: brewer’s yeast, glucose, sucrose, yeast extract, peptone, agar, MgSO₄ x 6H₂O, CaCl₂ x 2H₂O

Table 1. Propionic Acid Quantity and Approximate Yield/Container

Cat#	H ₂ O	Propionic Acid ^{††}	Narrow Yield	Wide Yield	Bottle Yield
66-115	1L	6.0ml	~110 Vials	~88 Vials	~22

^{††} Use suggested amount of propionic acid in Table 1 if using ≥99% (13.4M) propionic acid solution

Table 2. Suggested Volume/Drosophila Container

Volume	Container Type	Genesee Scientific Standard Vessel Catalog Numbers
~10ml	Narrow Vial	Polystyrene: #32-116 (bulk); 32-109 (tray pack)
~12ml	Wide Vial	Polystyrene: #32-117 (bulk); 32-110 (tray pack)
~50ml	Standard Bottle	Polypropylene: #32-130 / Glass: #50-102

Helpful Hints

- To remove clumps use a hand mixer, or press out larger lumps with fingers or the back of a large spoon.
- Make sure to let the food cool to between 60°C to 40°C before pouring. The medium will separate if it is too hot when poured.
- Problematic food is often simply dry food. Seemingly small changes in the cooking process can make a significant difference in the quality of the food if those changes result in increased water loss. Take special care in this regard, especially in dry climates.
- If new to Nutri-Fly start small on your first batches. Before combining multiple packets, start with one to gain experience and a comfort level and gradually move to larger batches.
- Even when hot the food is thick. Be sure your pump can handle it before you make a large batch.

^{††} Tegosept (Genesee #: 20-258; 20-259; 20-266 / synonyms: methyl paraben; nipagen) may be used in addition to but not instead of propionic acid. Propionic acid is considered sufficient as a preservative and the use of tegosept is known to slow Drosophila growth rates in the larval and pupal stages and we therefore suggest it be used only when additional fungal protection is necessary, or to eliminate sporadic mold problems.